



# Wissenschaftliches Rechnen II/Scientific Computing II

Sommersemester 2016  
Prof. Dr. Jochen Garcke  
Dipl.-Math. Sebastian Mayer



## Exercise sheet 11

To be handed in on **Thursday, 07.06.2016**

# Application of PCA and MDS

## 1 Group exercises

### G 1. (MDS: embedding of out-of-sample data)

Assume you had training data in the form of a centered Gram matrix  $G^c = (\langle y^i, y^j \rangle)_{i,j=1}^n = Y^T Y$  or in the form of a Euclidean distance matrix  $D = (\|y^i - y^j\|)_{i,j=1}^n$  and learned a  $p$ -dimensional embedding of the training data using the CMDS algorithm. Now assume there is a new test point  $x \in \mathbb{R}^d$ , which is different from the  $y^i$  but stems from the same data generating source. You cannot observe  $x$  directly but only one of the following sets of features:

- you either observe inner products  $x_S = Y^T x$ ,
- or you observe squared Euclidean distances  $x_E = (\|x - y^i\|_2^2)_{i=1}^n$ .

Use the components computed by the CMDS algorithm to construct a  $p$ -dimensional embedding  $\hat{x}$  of  $x$  from the given feature representation. Give a geometric interpretation of the constructed embedding  $\hat{x}$ . Discuss what properties the training data  $y^1, \dots, y^n$  must have such that the obtain embedding  $\hat{x}$  is reasonable.

*Solution.* I give a more detailed explanation, since some of you still struggled with how to interpret the embeddings generated by MDS/PCA. Let us first recapitulate how the training data is embedded. Consider the singular value decomposition  $Y = U \Sigma V^T$ . We have seen in the lecture that CMDS orthogonally projects the data  $Y$  on the  $p$ -dimensional subspace spanned by the columns of  $W = U I_{d \times p}$ , where  $W$  is such that the sum of squared errors (lengths of orthogonal complements)

$$\|Y - W W^T Y\|_F^2 = \sum_{i=1}^n \|y^i - W W^T y^i\|_2^2$$

becomes minimal. The embedding of the training data is given by  $W^T Y = I_{p \times d} U^T Y = I_{p \times d} \Sigma V^T$ . Note that  $\Sigma = \Lambda^{1/2}$ , where  $Y^T Y = V \Lambda V^T$ .

It seems natural to take

$$\hat{x} = W^T x = I_{p \times d} U^T x$$

as the embedding of the new data point  $x$ . How can we justify this? To this end, let us discuss another geometric interpretation of  $U$  and  $W$ , which can be considered as the non-probabilistic version of the PCA-perspective. The training data points  $y^1, \dots, y^n$  lie in an ellipsoid  $E_Y$ , which is given by the transformation of the the Euclidean unit ball  $B_2^d = \{y \in \mathbb{R}^d : \|y\|_2 \leq 1\}$  under the matrix  $Y Y^T$ ,  $E_Y = Y Y^T B_2^d$ . Now the nonzero eigenvalues on the diagonal of  $\Lambda$  and the columns of  $U$  describe the *principal axes* of this

ellipsoid. Concretely, the  $i$ th column of  $U$  contains the unit vector that is aligned with the  $i$ th principal axis of  $E_Y$  and the  $i$ th eigenvalues  $\lambda_i$  gives the length of this principal axis.

If the given training data are representative for the data generating source, future data points will also lie in  $E_Y$  (at least, they will be close to it). Hence, if we assume that the given data  $y^1, \dots, y^n$  is representative, then  $W^T x = I_{p \times d} U^T x$  will provide a good  $p$ -dimensional approximation of  $x$  in the sense that

$$\|x - WW^T x\|_2$$

is of the same magnitude as for the training data.

Now we are only left with the problem how to compute  $I_{p \times d} U^T x$  as we only know  $Y^T x$ . But this is simple: since  $Y^T = V \Sigma U^T$  we have  $U^T = \Sigma^{-1} V^T Y^T$ , such that

$$\hat{x} = I_{p \times d} \Sigma^{-1} V^T Y^T x.$$

In case that we only know Euclidean distances, then we use the same double centering trick which you have seen in the lecture:

$$\langle y^i, x \rangle = -\frac{1}{2} \left( \|y^i - x\|_2^2 - \frac{1}{n} \sum_{j=1}^n \|y^i - y^j\|_2^2 - \frac{1}{n} \sum_{j=1}^n \|x - y^j\|_2^2 + \frac{1}{n^2} \sum_{k,l=1}^n \|y^l - y^k\|_2^2 \right).$$

I leave it to you to write this in matrix form. □

## G 2. (Kernel-MDS)

Discuss how MDS could be generalized to distances and inner products which are induced by a reproducing kernel  $k : \Omega \times \Omega \rightarrow \mathbb{R}$ . Concretely, assume that there are points  $x_1, \dots, x_n \in \Omega$  of which you observe  $(k(x_i, x_j))_{i,j=1}^n$  and you want to construct embeddings  $\hat{x}^1, \dots, \hat{x}^n \in \mathbb{R}^p$  such that

$$k(x_i, x_j) \approx \langle \hat{x}^i, \hat{x}^j \rangle.$$

*Solution.* Consider the eigenvalue decomposition of  $K$ ,  $K = V \Lambda V^T$  or  $\Lambda = V^T K V$ . Let us denote by  $V_{\cdot j}$  the  $j$ th column of  $V$ . Since  $k(x_i, x_j) = \langle k(x_i, \cdot), k(x_j, \cdot) \rangle_k$ , we have

$$\lambda_i \delta_{ij} = V_{\cdot i}^T K V_{\cdot j} = \sum_{l_1, l_2=1}^n V_{l_1 i} k(x_{l_1}, x_{l_2}) V_{l_2 j} = \left\langle \sum_{l=1}^n V_{li} k(x_l, \cdot), \sum_{l=1}^n V_{lj} k(x_l, \cdot) \right\rangle_k,$$

where  $\delta_{ij}$  is as usual the Kronecker delta. Assuming that  $K$  has rank  $n$ , the functions

$$f_i = \frac{1}{\sqrt{\lambda_i}} \sum_{l=1}^n V_{li} k(x_l, \cdot), \quad i = 1, \dots, n$$

are pairwise orthogonal and  $\|f_i\|_k = 1$ .

To understand what these functions mean, consider the *feature map*

$$\Phi : \Omega \rightarrow H, \quad x \mapsto k(x, \cdot).$$

Then  $\Phi$  maps the given data points  $X = \{x_1, \dots, x_n\}$  into a  $n$ -dimensional subspace of  $\mathcal{H}$ , namely  $\mathcal{H}_X = \text{span}\{k(x_i, \cdot), i = 1, \dots, n\}$ . Since  $f_i \in \mathcal{H}_X$ ,  $(f_i)_{i=1}^n$  is an orthonormal basis of  $\mathcal{H}_X$ . As you will show in H1, this ONB has the property that  $f_1, \dots, f_p$  span the optimal  $p$ -dimensional subspace of  $\mathcal{H}_X$  such that

$$\sum_{i=1}^n \|k(x_i, \cdot) - \sum_{j=1}^p \langle k(x_i, \cdot), f_j \rangle f_j\|_k^2$$

becomes minimal. Note that  $\langle k(x_i, \cdot), f_j \rangle = f_j(x_i)$ . Hence, for sufficiently large  $p$ ,

$$k(x_i, x_j) \approx \left\langle \sum_{l=1}^p \langle k(x_i, \cdot), f_l \rangle f_l, \sum_{l=1}^p \langle k(x_j, \cdot), f_l \rangle f_l \right\rangle = \sum_{l=1}^p f_l(x_i) f_l(x_j).$$

Thus, we choose as embedding

$$\hat{x}^i = \begin{pmatrix} f_1(x_i) \\ \vdots \\ f_p(x_i) \end{pmatrix}.$$

□

## 2 Homework

**H 1.** (Optimal  $p$ -dimensional subspace in a RKHS)

Let  $k : \Omega \times \Omega \rightarrow \mathbb{R}$  be a reproducing kernel,  $\mathcal{H}$  its native Hilbert space and  $X = \{x_1, \dots, x_n\} \subset \Omega$ . Consider the kernel matrix  $K = (k(x_i, x_j))_{i,j=1}^n$  and the corresponding eigenvalue decomposition  $K = V\Lambda V^T$  with  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ , where we assume  $m \leq n$ . Consider for  $i = 1, \dots, m$  the functions

$$f_i := \frac{1}{\sqrt{\lambda_i}} \sum_{j=1}^n V_{ji} k(x_j, \cdot) \in \mathcal{H}_X.$$

a) Show that  $(f_i)_{i=1}^m$  forms an orthonormal basis of  $H_X$ . **Hint:** Use that  $\Lambda = V^T K V$ .

b) Let  $p \in \{1, \dots, m\}$ . Show that  $(f_i)_{i=1}^p$  is the solution of

$$\min_{(g_i)_{i=1}^p} \min_{\text{ONB of } \mathcal{H}_X} \sum_{i=1}^n \|k(x_i, \cdot) - \sum_{j=1}^p g_j(x_i) g_j\|_k^2.$$

**Hint:** Argue analogously as in Sheet 10, H2 b).

(10 Punkte)

*Solution.*

a) Orthogonality has already been shown in G2, see the solution there. It only remains to note that since  $K$  has rank  $m$ , the subspace  $\mathcal{H}_X$  is  $m$ -dimensional and hence the  $f_i$  form an ONB.

b) Let us first observe, using the symmetry of  $k$  and  $\Lambda = V^T K V$ ,

$$\begin{aligned} \sum_{l=1}^n f_i(x_l) f_j(x_l) &= \frac{1}{\sqrt{\lambda_i \lambda_j}} \sum_{t_1, t_2=1}^n V_{t_1 i} V_{t_2 j} \sum_{l=1}^n k(x_{t_1}, x_l) k(x_l, x_{t_2}) \\ &= \sum_{t_1, t_2=1}^n V_{t_1 i} V_{t_2 j} (K^2)_{t_1 t_2} = V_{\cdot i}^T K^2 V_{\cdot j} = \sqrt{\lambda_i \lambda_j} V_{\cdot i}^T V_{\cdot j} = \lambda_i \delta_{ij}. \end{aligned}$$

Now, using the orthogonality of the  $g_j$ , we compute

$$\sum_{i=1}^n \|k(x_i, \cdot) - \sum_{j=1}^p g_j(x_i) g_j\|_k^2 = \sum_{i=1}^n k(x_i, x_i) - \sum_{i=1}^n \sum_{j=1}^p g_j(x_i)^2.$$

This sum becomes minimal when  $\sum_{i=1}^n \sum_{j=1}^p g_j(x_i)^2$  is maximal. Since the  $f_i$  form an ONB, we have  $k(x_i, \cdot) = \sum_{l=1}^m f_l(x_i) f_l$ , which yields

$$\sum_{i=1}^n \sum_{j=1}^p g_j(x_i)^2 = \sum_{i=1}^n \sum_{j=1}^p |\langle g_j, k(x_i, \cdot) \rangle_l|^2 = \sum_{i=1}^n \sum_{j=1}^p \sum_{l_1, l_2=1}^m \langle g_j, f_{l_1} \rangle_k \langle g_j, f_{l_2} \rangle_k f_{l_1}(x_i) f_{l_2}(x_i)$$

Our first observation implies

$$\sum_{i=1}^n \sum_{j=1}^p \sum_{l_1, l_2=1}^m \langle g_j, f_{l_1} \rangle_k \langle g_j, f_{l_2} \rangle_k f_{l_1}(x_i) f_{l_2}(x_i) = \sum_{j=1}^p \sum_{l=1}^m |\langle g_j, f_l \rangle_k|^2 \lambda_l$$

Put  $\alpha_l = \sum_{j=1}^p |\langle g_j, f_l \rangle_k|^2$ . By Bessel's inequality,  $\alpha_l \leq \|f_l\|_k^2 = \lambda_l$ . Since  $g_j = \sum_{l=1}^m \langle g_j, f_l \rangle_k f_l$ , we further have

$$\sum_{l=1}^m \alpha_l = \sum_{l=1}^m \sum_{j=1}^p \|\langle g_j, f_l \rangle_k f_l\|_k^2 = \sum_{j=1}^p \left\| \sum_{l=1}^m \langle g_j, f_l \rangle_k f_l \right\|_k^2 = p.$$

It remains to apply Sheet 10 G2 to conclude

$$\sum_{i=1}^n \sum_{j=1}^p g_j(x_i)^2 = \sum_{l=1}^m \alpha_l \lambda_l \leq \sum_{l=1}^p \lambda_l.$$

We have equality if  $g_i = f_i$  for  $i = 1, \dots, p$ .

□

**H 2.** (Programming exercise: pedestrian classification)

See accompanying notebook.

(10 Punkte)