



Wissenschaftliches Rechnen II/Scientific Computing II

Sommersemester 2016
Prof. Dr. Jochen Garcke
Dipl.-Math. Sebastian Mayer



Exercise sheet 12

To be handed in on **Thursday, 14.07.2016**

Isomap

1 Group exercises

G 1. Consider the manifold $M = \{x \in \mathbb{R}^d : \|x\|_2 = r\}$, i.e., the Euclidean sphere with radius $r > 0$ in \mathbb{R}^d . Compute the minimum radius of curvature $r_0(M)$ and the minimum branch separation $s_0(M)$. Moreover, show that Lemma 2.16 holds true with equality, i.e., for all $x, y \in M$ such that $d_M(x, y) < \pi r_0(M)$ we have

$$d_E(x, y) = 2r_0(M) \sin\left(\frac{d_M(x, y)}{2r_0(M)}\right).$$

G 2. Construct a graph distance matrix which is not a Euclidean distance matrix.

G 3. Let $M \subset \mathbb{R}^d$ be manifold. Assume you have run Isomap with input $x_1, \dots, x_n \in M$ to obtain p -dimensional embeddings. Write down ready-to-use formulas that compute you the p -dimensional embedding for a new, unseen data point $x \in M$ without rerunning the whole Isomap algorithm.

2 Homework

H 1. Let M be a compact manifold. Prove the following simplified version of Lemma 2.16. For any $\varepsilon > 0$, we have

$$(1 - \varepsilon)d_M(x, y) \leq d_E(x, y) \leq d_M(x, y)$$

for all $x, y \in M$ such that $d_M(x, y) < 2r_0(M)\varepsilon$. **Hint:** Consider a unit speed parametrization $\gamma : (0, l) \rightarrow M$ with $\gamma(0) = x$, $\gamma(l) = y$ and use the fundamental theorem of calculus to obtain a first order estimate.

(10 Punkte)

H 2. (Isomap and non-EDM graph distance matrices)

a) Let S be a symmetric $n \times n$ matrix with eigenvalue decomposition $U\Lambda U^T$. Let Λ' be Λ with all negative eigenvalues replaced by zero and put $S^+ := U\Lambda'U^T$. Show that S^+ is the solution of

$$\min_{B \in \mathcal{S}_n^+} \|S - B\|_F^2,$$

where \mathcal{S}_n^+ is the set of all positive semi-definite $n \times n$ matrices.

b) Based on your insights from a), argue why Isomap can also be used when the graph distance matrix D_G is not a Euclidean distance matrix. How should the algorithm be modified in this case?

(10 Punkte)